The Road Traveled to "No Elective Deliveries less than 39 Completed Weeks"

By Colette B. LaCombe, RNC



I wish to thank Donna Frye, RN, MN, Director, HCA Women's & Children's Clinical Service for providing the information on behalf of HCA.

WHO ARE WE

OUR ORGANIZATION



Central Group Mid America Division





A Campus of THE REGIONAL MEDICAL CENTER OF ACADIANA

Women's & Children's Hospital Lafayette, LA





A Campus of THE REGIONAL MEDICAL CENTER OF ACADIANA

Women's & Children's Hospital

Founded in 1983, Women's & Children's Hospital, A campus of The Regional Medical Center of Acadiana has been providing specialized health care services for women and children for more than 20 years.

Since our humble beginnings as a birthing hospital, we have grown to become one of Louisiana's leading health care institutions dedicated exclusively to the care of women and children.



Women's & Children's Hospital

Awards/Accreditation:

- TJC Lab Accreditation
- CLIA Accreditation
- NICU "Center of Excellence" designation from United Healthcare
- Voted "Best Place to have a Baby" Times of Acadiana"
- Ambulatory Surgery Center Lafayette Surgicare
- Critical Care
 - NICU; NICU Transport
 - PICU; PICU Transport
 - Pediatric Emergency Room
- Diagnostic Imaging Services
- Patient Education and Support groups
 Emergency Services

- Laboratory
- Level III Neonatal Intensive Care Unit (Regional Referral Center)
- Pediatrics
- Pediatrics Specialty Clinic
- Pediatric Surgical Specialties
- Sleep Lab
- Surgery
- Women's Services including
 - Dedicated High Risk Antenatal Unit
 - Fertility
 - Labor/Delivery/Recovery Suites
 - OB Hospitalist
 - Postpartum Unit



THE REGIONAL MEDICAL CENTER OF ACADIANA

LABOR & DELIVERY / HIGH RISK ANTENATAL UNIT

- Labor and Delivery Unit
 - 2010 yearly deliveries 3245
 - 15 LDRs
 - 2 Bed OB Triage
 - 2 OB OR Suites
 - 2 Bed OB Recovery
- High Risk Antenatal Unit
 - 10 Beds
 - Maternal Fetal Medicine Program

HCA STORY

The annual 2006 HCA Perinatal Conference focused on care of the "near term premature infant".

Discussions raised questions --

- Were there totally elective deliveries and cesarean sections? What was the frequency?
- Is iatrogenic prematurity a problem?
- Are there implications for the infants born at term, less than 39 week?
- Can we improve outcomes by minimizing elective deliveries prior to 39 weeks gestation?
- What evidence does the evidence say?

Our Focus became the 37-39 week elective induction.

2007 STUDY

2007 STUDY OBJECTIVES

- To determine the frequency of elective term deliveries prior to 39 completed weeks in the United States population.
- To examine neonatal outcomes associated with elective delivery at "term" but prior to 39 completed weeks of gestation in all forms of deliveries (inductions, primary and repeat cesarean sections).
- To examine maternal consequences of elective term inductions.

STUDY METHODOLOGY

- Hospital Corporation of America 114 obstetric facilities in 21 states.
- ▶ 225, 000 annual deliveries.
- Has been shown to be representative of the U.S. population as a whole.
 - Clark, S., Belfort, M., Dildy, G., Herbst, M., Meyers, J., & Hankins, G. (2008). Maternal death in the 21st century: Causes, prevention, and relationship to cesarean delivery. American Journal of Obstetrics and Gynecology. In Press.
 - Clark, S. L., Belfort, M. A., Hankins, G. D., Meyers, J. A., & Houser, F. M. (2007).
 Variation in the rates of operative delivery in the United States. *American Journal of Obstetrics and Gynecology*, 196(6), 526.e1-526.e5.

2007 STUDY METHODOLOGY

- Population sampled: All deliveries between May 1, 2007 and July 31, 2007 in 27 HCA facilities.
 - Women's & Children's Hospital
- Facilities were selected to be representative of the entire population—geographic characteristics and delivery volume –
- Comprehensive data was collected for all women undergoing planned delivery at 37 weeks and 0 days or greater.

2007 STUDY RESULTS

- ▶ 17, 794 deliveries
- ▶ 14,955 at 37 weeks or greater
- 6562 were planned term deliveries.
- ▶ 4645 were <u>elective planned term deliveries.</u>
- > 31% of all term deliveries were elective.
- 11% of all term deliveries were elective and prior to 39 completed weeks gestation.
 - THIS IS WHERE WE CAN MAKE A DIFFERENCE.

RESULTS NICU Admissions Following *Elective*Deliveries

- ▶ 37.0 37.6 weeks: 17.8%
- ▶ 38.0 38.6 weeks: 8.2%
- > 39 weeks: 4.6%
- All differences highly significant (p<0.001)</p>
- 2/3 were direct NICU admits, 1/3 were admitted later after initial normal newborn admission.
 - As a note, the delivery provider may not realize the baby went to the NICU after the initial admission.
- Mean NICU stay for these infants was 4.5 days.

2007 STUDY CONCLUSIONS

- ▶ 11% of all term deliveries are elective and performed prior to 39 weeks gestation, contrary to longstanding ACOG/AAP recommendations.
- Given the nature of many "indications", the actual rate is probably higher.
- Such infants experience significant morbidity.
- For all Planned Inductions, the cesarean delivery rate is directly related to initial cervical dilatation.
- Elective Induction of labor with an unfavorable cervix also increases the risk of cesarean delivery.

DISSEMINATION OF THE FINDINGS

• Fall 2007

- Findings were shared with the Clinical Work Group
- Findings were shared at the Perinatal Conference
- Some Hospitals implemented policies

Spring 2008

 Supported a National Quality Form Task Force to establish National Voluntary Consensus Standards for Perinatal Care

ADOPTION AND IMPLEMENTATION

Fall of 2008

Tool boxes including references, educational resources, and a prototype policy were provides to the hospitals

Dr. Clark conducted "Physician to Physician" Conference calls

Monitoring of Perinatal Services to determine:

Discussion

Adoption of a position statement Endorsement of a policy

ADOPTION AND IMPLEMENTATION

Additional 2008 Strategies

Perinatal nursing leaders provided education to physician office staff.

Perinatal nursing leaders, childbirth educators, and staff nurses provided targeted patient *education—correcting misconceptions regarding the safety of early term births*.

Physician champions and Nursing leaders began to provide education to OB department s and advocate for a position statement or a policy regard "No Elective Deliveries Less than 39 weeks gestation."

ELECTIVE DELIVERIES PRIOR TO 39 WEEKS

2009

The Joint Commission adopted as a core measure

Leapfrog has adopted as a quality measure May

76 of the 111 hospitals or 68% have adopted a position statement or endorsed a policy Replication of the study began April Data collection for compliance began

2009 STUDY

OBJECTIVE AND STUDY DESIGN

Objective:

Examine the effectiveness of different approaches to a reduction in elective early term deliveries or the effect of such policies on newborn intensive care admissions and still birth rates.

Study Design:

We conducted a retrospective cohort study of prospectively collected data and examined outcomes in 27 hospitals before and after implementation of 1 of 3 strategies for the reduction of elective early term deliveries.

THREE APPROACHES TO PRACTICE CHANGE:

All hospitals began with similar rates of elective delivery at <39 weeks of gestation.

Group 1

Endorsed a formal policy enforced by hospital staff.

Group 2

Endorsed a formal policy that was not enforced by hospital staff, but with automatic peer review for exceptions.

Group 3


Relied on education only.

2009 STUDY RESULTS:


| Variable | 2007 | 2009 | P value |
|--|-----------------|---------------|---------|
| Deliveries, n | 17,194 | 17,221 | NA |
| Deliveries > or = 37 wk, n | 14,995 | 14,863 | NA |
| Planned +Elective Deliveries at 37.0-38.6 wk, n | 6,562 | 4349 | <.001 |
| Elective Deliveries at 37/0-38.6 wk, n (%) | 1712 (9%) | 746 (4.3) | <.001 |
| Group 1:7 hospitals, n/N(%) | 320/3886 (8.2) | 5/3818 (1.7) | .007* |
| Group 2:9 hospitals, n/N(%) | 403/4797 (8.4) | 155/4646(3.3) | <.025 |
| Group3:11 hospitals, n/N(%) | 989.9111 (10.9) | 526/8757(6.0 | .135* |
| Neonatal Intensive Care Unit admissions at $> or=37$ wk, n (%) | 1328 (8.9) | 1119 (7.5) | <.001 |

^{*}indicates 2-way analysis of variance, other P values were obtained with X2 Yates correlation correction

TERM DELIVERIES 2007 VS 2009 Facilities



% Reduction in elective birth < 39 weeks, according to policy approach



■ % reduction from 2007baseline

2009 STUDY RESULTS AND CONCLUSIONS

- Elective early term delivery was reduced from 9.6-4.3% of deliveries.
- The rate of term neonatal intensive care admissions fell by 16%.
- We observed no increase in stillbirth.
- The greatest improvement was seen when elective deliveries at <39 weeks were not allowed by hospital personnel.</p>
- Physician education and the adoption of policies backed only by peer review are less effective than "hard stop" hospital policies to prevent elective deliveries prior to 39 weeks gestation.

2009 STUDY RESULTS AND CONCLUSIONS

- Elective early term delivery was reduced from 9.6-4.3% of deliveries.
- The rate of term neonatal intensive care admissions fell by 16%.
- We observed no increase in stillbirth.
- The greatest improvement was seen when elective deliveries at <39 weeks were not allowed by hospital personnel.</p>
- Physician education and the adoption of policies backed only by peer review are less effective than "hard stop" hospital policies to prevent elective deliveries prior to 39 weeks gestation.

WOMEN'S & CHILDREN'S HOSPITAL JOURNEY

WOMEN'S & CHILDREN'S HOSPITAL JOURNEY

- ▶ In 2007, Participated in the 1st study
- In Fall of 2008, adopted a soft approach
 - Physician & Staff education conducted
- In Early 2009, received the HCA Tool Box
- Embarked on our journey towards reducing elective deliveries less than 39 completed weeks
- ▶ In May 2009, participated in the 2nd study

WOMEN'S & CHILDREN'S HOSPITAL JOURNEY

- July 2009, Physicians agree to adopt policy
 Policy adopted however no penalties for scheduling
- In 2010 & 2011, policy revised to include consequences for policy violation
- In February 2011, WCH approached to join state-wide collaborative
- March 2011, enrolled in IHI Project

The Team

- Colette LaCombe, RNC Team Leader
- Dr. Frank Caillet, Chief of OB Services
- Dr. Sheryl Rodts-Palenik, MFM
- Sharon Gates, RN, ACNO
- John Marker, CNO
- Bill Ferry, Director, Risk Management
- Ginger Broussard, Director, Quality
- Anitra Ford-Scott, RN, L & D Manager
- Paige Landry, RN, Staff
- Melanie Romero, RNC, Staff

IHI PROJECT MEASURE SETS

- Elective Delivery Rate prior to 39 weeks
- Cesarean Rate for low risk first birth women
- Perinatal Harm
- Gestational Age Reliability
- Transfer to Higher Level Neonate

IHI Project

- First Steps
 - Team
 - What are we suppose to be doing?
- Data collection
- Challenges
 - Physicians
 - Patients
 - Scheduling
- Next Steps

PATIENTS DELIVERED BETWEEN > 37 WEEKS AND < 39 WEEKS DATA REPORTED

Percentage of patients who electively delivered newborns with >= 37 and < 39 weeks of gestation completed (TJC PC.01)

| Lafayette Women's and Children's Hospital – Louisiana | | | | | |
|---|---------------|--------------------------------------|---|-----------------|--|
| Time Period | Percent Value | Patients with Elective Deliveries | Patients delivering >=37 and <39 weeks gestation | Annotation Type | |
| 5 - 2011 | 20.19 | 21 | 104 | None | |
| 6 - 2011 | 13.89 | 10 | 72 | None | |

Next Steps

- Revise the policy with a "Hard Stop"
- Revised scheduling procedure for inductions / C-Sections
- Revise forms

REFERENCES:

- Alan TN, Landon Mark, Spong CY et al: Timing of Elective Repeat Cesarean Delivery at Term and Neonatal Outcomes. New England Journal of Medicine, January 2009
- American College of Obstetricians and Gynecologists. ACOG Practice Bulletin no. 107: Induction of Labor. Obstetrics and Gynecology 2009; 114: 386-97.
- American College of Obstetricians and Gynecologist Technical Bulletin no 97.
 Fetal Lung Maturity. September 2008
- American College of Obstetricians and Gynecologists. Late Preterm Infants. Washington, DC: The College: 2008. Committee Opinion no. 404.
- Clark SL, Belfort MA, Miller DK et al: Neonatal and Maternal Outcomes associated with elective term delivery. American Journal of Obstetrics and Gynecology, January 2009
- Clark S, Frye D, Meyers J, Belfort M, Dildy G, Kofford S, Englebright J, Perlin J. Reduction in elective delivery at elective 39 weeks of gestation: comparative effectiveness of 3 approaches to change and the impact on neonatal intensive care admission and stillbirth. American Journal of Obstetrics and Gynecology 2010: 1.e1-6.

QUESTIONS?

THANK YOU!